Stepan_®



Product Bulletin

Product Name

Chemical Structure

CAS Registry No. **INCI Name**

Applications

Typical **Properties**

Biodegradability

Toxicity

Storage & Handling

Clearances

AMMONYX® LO

$$CH_{3}$$

$$CH_{3}(CH_{2})_{X} - N \rightarrow O$$

$$CH_{3}$$

$$X = 11 \text{ (predominantly)}$$

1643-20-5

Lauramine Oxide

Functional Properties

- Foam Enhancer
- Foam Stabilizer
- Viscosity Builder
- Lubricant*
- Bleach Stable

(*For the textile industry)

Emulsifier*

- Wetting Agent*
- Dye Dispersant*
- Can be formulated with anionic, nonionic and cationic materials.

Acid Stable

End Product Uses

Personal Care

- Shampoos
- Handsoaps
- **Bath Products**
- Creams & Lotions
- Hair Conditioners

Household & I&I

- **Dishwashing Detergents**
- Cleansers
- Hard Surface Cleaners (which contain acids or bleach)

Appearance at 25 °C Clear liquid	Color (APHA), as is 100 max.
Amine Oxide (MW 251), %	Viscosity at 25 °C, cps
Free Amine (MW 235), % 0.49	Flash Point (PMCC), °C (°F) >94 (>201)
pH, 10% aqueous 8.0	Density, g/ml (lbs/U.S. gal) 0.968 (8.05)
Cloud Point (as is), °C (°F)1 (30)	RVOC, U.S. EPA, % 0
Boiling Point, °C (°F) 100 (212)	Free Peroxide, % 0.13
Preservative	

Product is readily biodegradable. A detailed biodegradability statement is available upon request.

The available toxicity data shows AMMONYX LO to be slightly toxic orally to rats, (LD₅₀ = 3600 mg/kg). At 2.5% active, AMMONYX LO may cause severe eye irritation and minimal irritation to the skin.

Normal safety precautions (i.e., gloves and safety goggles) should be employed when handling AMMONYX LO. Contact with the eyes and prolonged contact with the skin should be avoided. Wash thoroughly after handling material.

It is recommended that AMMONYX LO be stored in sealed containers and kept at temperatures not lower than 50 °F (10 °C) nor higher than 120 °F (49 °C). Avoid overheating or freezing.

Bulk Storage Information: Tanks, with a conservation vent, made of 316L stainless steel or fiberglass with an Atlac 382 corrosion liner are recommended. Centrifugal or positive displacement gear pumps and piping should be 316L stainless steel. Recommended storage for bulk tanks is 20-52 °C (68-125 °F).

Standard Packaging: AMMONYX LO is available in bulk and in 55 gallon drums (net weight 425 lb/193 kg).

All components of AMMONYX LO are listed in the following countries; the registration numbers for the active ingredients are included in parentheses: United States (TSCA 1643-20-5), Europe (EINECS 216-700-6), Japan (ENCS 2-198), Canada (DSL 1643-20-5), Australia (AICS 1643-20-5), Korea (ECL Serial No. KE-11348), Philippines (PICCS 1643-20-5) and China (EICS 1643-20-5).

AMMONYX LO is cleared for use user 40 CFR 180.910, pre- and post-harvest applications. AMMONYX LO meets the U.S. EPA Safe Choice Program's surfactant screen and is listed on www.cleangredients.org (an on-line

Oct. 2017 Supersedes; March 2013 Page 1 of 2



Formulations

database for green formulators). For further information, visit www.cleangredients.org. Additional Stepan ingredient listing can be found at www.stepan.com.

CONDITIONING SHAMPOO

<u>Ingredients</u>		Wt, % (as is)	<u>Function</u>	
1.	AMPHOSOL® CA	32.8	Cosurfactant	
2.	AMMONYX LO	32.0	Cosurfactant	
3.	AMMONYX 4	1.7	Conditioning Agent	
4.	Preservative, Dye, Fragrance	q.s.	Additives	
5.	Citric Acid (50%)	q.s.	pH Adjuster	
6.	Distilled Water	q.s. to 100.0	Solvent, Carrier	
7.	Sodium Chloride	q.s.	Thickener	

Mixing Procedure:

Add AMPHOSOL CA, AMMONYX LO and AMMONYX 4 to water. Heat with mixing to 40-50 °C (104-122 °F), until AMMONYX 4 is completely blended into solution. Cool to 30-35°C (86-95°F). Add perfume, preservative and dye, as desired. Adjust pH with citric acid to 6.0 - 6.5. Add sodium chloride to achieve desired viscosity. Changing pH changes properties.

Physical Properties:

pH (as is)	
Appearance at 25 °CLi	iauid
GENTLE SHAMPOO	•

		GENTLE SHAWFOO	
Ingred	<u>lients</u>	Wt, % (as is)	<u>Function</u>
1.	STEOL® CS-330	30.0	Primary Surfactant
2.	STEPANOL [®] AM-V	15.0	Secondary Surfactant
3.	AMMONYX LO	1.0	Foam Enhancer/Foam Stabilizer
4.	NINOL® 40-CO	2.0	Foam Enhancer/Viscosity Booster
5.	Glycerin	3.0	Humectant
6.	Methocel E4M (Dow Chemical)	0.4	Thickener
7.	Sodium Chloride	1.0	Thickener
8.	Citric Acid (50%)	q.s.	pH Adjuster
9.	Preservative, Dye, Fragrance	q.s.	Additives
10.	Distilled Water	q.s. to 100.0	Solvent, Carrier

Mixing Procedure:

Heat one third of the required water to 80 °C (176 °F). Add Methocel E4M and agitate until all particles have been dispersed. Turn off heat and add the remainder of the water. Agitate until the solution is clear and particle free. Add items 1 through 6 and blend until homogeneous. Adjust pH with citric acid to 6.0 - 6.5. Viscosity may be modified by adjusting the sodium chloride content.

Physical Properties:

pH (as is)	6.0 - 6.5
Appearance at 25 °C	Liquid

This product bulletin has been written in accordance with ACC's Product Stewardship guidelines.

A Safety Data Sheet is available upon request.



The information contained herein is based on the manufacturer's own study and the works of others and is subject to change without prior notice. The information is not intended to be all-inclusive, including as to the manner and conditions of use, handling, storage or disposal or other factors that may involve additional legal, environmental, safety or performance considerations. Nothing contained herein grants or extends a license, express or implied, in connection with any patents issued or pending of the manufacturer or others, or shall be construed as a recommendation to infringe any patents. STEPAN COMPANY MAKES NO PRODUCT WARRANTY OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR USE, EXPRESS OR IMPLIED, AND NO OTHER WARRANTY OR GUARANTY, EXPRESS OR IMPLIED, IS MADE, INCLUDING AS TO INFORMATION REGARDING PERFORMANCE, SAFETY, SUITABILITY, STABILITY, ACCURACY, COMPLETENESS, OR ADEQUACY. Stepan Company (and its employees, subsidiaries and affiliates) shall not be liable (regardless of fault) to the vendee, its employees, or any other party for any direct, indirect, special or consequential damages arising out of or in connection with the accuracy, completeness, adequacy, furnishing, use, or reliance upon information provided herein. The vendee assumes and releases Stepan Company (and its employees, subsidiaries and affiliates) from all liability, whether in tort, contract or otherwise to the fullest extent possible under the relevant law."

tepan

Corporate Headquarters Northfield, Illinois 60093, U.S.A.

Phone: 847-446-7500 Fax: 874-501-2100

Website: www.stepan.com

For Technical Service Call: Stepan North America +1-800-745-7837 Stepan Mexico +55-11-5089-2208 Stepan Quimica +55-11-5089-2200 Stepan Colombia +571-636-2808

Stepan Europe +33-476-505-160 Stepan Asia +656-879-9855 Stepan Philippines +656-879-9855 For Customer Service Call:

Stepan North America +1-800-457-7673 Stepan Mexico +52-555-533-1967 Stepan Quimica +55-11-5089-2200 Stepan Colombia +57-6-8993125 Stepan UK +44-161-338-5511 Stepan France +33-476-50-51-00 Stepan Philippines +63-2-842-7951

Oct 2017 Supersedes; March 2013 Page 2 of 2

Product

Stewardship

Additional

Safety Information