Technical Information

Stepan

Stepan Company

Northfield, Illinois 60093 Telephone 847 446 7500



MAMMALIAN TOXICOLOGY OF DIMETHYL AMIDES

Applicable to these current Stepan products:

HALLCOMID™ 1025	STEPOSOL® MET-10U	
-----------------	-------------------	--

Toxicological Information:

Test/Conditions	Results/Classification	<u>References</u>
Acute Oral Toxicity (rat) (14 day) (gavage)	$LD_{50} = 550 \text{ mg/kg}$ (slightly toxicity)	Stepan Study No. 14-001A
Bovine Corneal Opacity	30.8	Stepan Study No. 13-019A
Permeability Assay	(moderate eye irritation)	
Skin Irritation Test (Epiderm model)	Mean tissue viability < 50% (irritating to skin)	Stepan Study No. 13-019B

References:

®; STEPOSOL® are registered trademarks of Stepan Company.

Last Update: 04/21/2016 Revision Reference: TX061-00

Nothing contained herein grants or extends a license, express or implied, in connection with patents, issued or pending, of the manufacturer or others. The information contained herein is based on the manufacturer's own study and the works of others. The manufacturer makes no warranties, expressed or implied, as to the accuracy, completeness, or adequacy of the information contained herein. The manufacturer shall not be liable (regardless of fault) to the vendee's employees, or anyone for any direct, special or consequential damages arising out of or in connection with the accuracy, completeness, adequacy or furnishing of such information.